



XIX CONGRESSO
NAZIONALE
SIES 2026

**FLT3 AS A NOVEL THERAPEUTIC TARGET IN PRECLINICAL
MODEL OF PEDIATRIC PAX5-REARRANGED BCP-ALL**

Alessia Curto, PhD Student

University of Milano-Bicocca, Fondazione Tettamanti

Firenze | 4-6 marzo 2026
Palazzo degli Affari




Disclosures of Alessia Curto

Not applicable

Introduction

- **B-cell Acute Lymphoblastic Leukemia (B-ALL)** is the most common pediatric malignancy
- **Characterized by molecular alterations** that define the different subgroups

- **PAX5-alt** is linked to poorer prognosis and a lower chance of survival and complete remission




Jia and Gu, Frontiers in Oncology 2022

C051

LA LEUCEMIA LINFOBLASTICA ACUTA (LLA) PEDIATRICA PAX5-R È UN SOTTOGRUPPO MOLECOLARE AD ALTO RISCHIO DI LLA A CELLULE B CARATTERIZZATO DA UN PROFILO PROGNOSTICO SFAVOREVOLE MRD-DIPENDENTE


N. Peccatori, A. Curto, D. Silvestri, S. Rebellato, Ž. Antić, S. Bhatia, A.K. Bergmann, A. Borkhardt, M. Zimmermann, S. Strehl, K. Nebral, C. Saitta, M. Quadri, C. Palmi, M. Bardini, D. Guardo, L. Lo Nigro, R. Parasole, M.C. Putti, F. Locatelli, J. Alten, M. Stanulla, M.G. Valsecchi, B. Buldini, V. Conter, A. Biondi, A. Balduzzi, M. Schrappe, A. Attarbaschi, G. Cario, C. Rizzari, G. Cazzaniga, G. Fazio (Monza-MB, Milano, Hannover-DE, Würzburg-DE, Düsseldorf-DE, Vienna-AT, Genova, Catania, Napoli, Padova, Roma, Kiel-DE)



Fazio G, Bresolin S, Silvestri D, et al. PAX5 fusion genes are frequent in poor risk childhood acute lymphoblastic leukaemia and can be targeted with BIBF1120

Hypothesis of the study


a.



b.



FLT3 gene




Study objectives

- Study of the *Flt3* gene expression profile in patients with *PAX5r* B-ALL to identify a potential innovative therapeutic target
- Evaluation of cell viability following treatment with FLT3 inhibitors *in-vitro*, *ex-vivo* e *in-vivo*

Study objectives

- Study of the *Flt3* gene expression profile in patients with *PAX5r* B-ALL to identify a potential innovative therapeutic target
- Evaluation of cell viability following treatment with FLT3 inhibitors *in-vitro*, *ex-vivo* e *in-vivo*

Results (I): high levels of *FLT3* expression in *PAX5r* patients



Study objectives

- Study of the *Flt3* gene expression profile in patients with *PAX5r* B-ALL to identify a potential innovative therapeutic target
- Evaluation of cell viability following treatment with FLT3 inhibitors *in-vitro*, *ex-vivo* e *in-vivo*

Results (II): HTP Drug screening in PAX5r PDXs

FLT3 expression levels from RNA-SEQ






Image adapted from Oikonomou A., Valsecchi L., Quadri M., et al. High-throughput screening as a drug repurposing strategy for poor outcome subgroups of pediatric B-cell precursor Acute Lymphoblastic Leukemia


Results (II): Gilteritinib has a potential new drug for PAX5r in pediatric B-ALL

FLT3 inhibitors on PAX5r




WHY GILTERITINIB?


- **Selective** for FLT3
- **Safe** for patients
- **FDA-approved** for adult patients with FLT3-mutated relapsed/refractory acute myeloid leukemia (AML) and in **pediatric patients** with relapsed/refractory FLT3-ITD AML



Results (II): Gilteritinib is efficient in *PAX5r* cell line



- NALL-1 (*PAX5::ETV6*) ←
- SEM (*KMT2A::AFF1*)
- ▲ 697 (*TCF3::PBX1*)
- ▼ RCH-ACV (*TCF3::PBX1*)









Cell line	IC50 Gilte (nM)
NALL-1	1014
SEM	112,4
697	7170
RCH-ACV	884,3

*The evaluation of gilteritinib efficacy in JIH-5 (*EP300::ZNF384*) is ongoing..




Results (II): Gilteritinib reduces cell viability in PAX5r PDX cells in a dose-dependent manner

Number of live cells (Ann V-/7-AAD-) in treated vs untreated condition
n=3 experiments




One-way ANOVA with the Dunnett post-test for multiple comparisons using untreated as the comparator






Results (II): Gilteritinib exerts a synergistic effect in combination with dexamethasone



Results (II): *in-vivo* experimental design




Results (II): gilteritinib demonstrates potent antileukemic activity *in-vivo*



Peripheral blood (PB) was also evaluated, but it showed poor engraftment (0%-0,4%)

One-way ANOVA with the Dunnett post-test for multiple comparisons using untreated as the comparator

Results (II): gilteritinib demonstrates potent antileukemic activity *in-vivo*



Peripheral blood (PB) was also evaluated, but it showed poor engraftment (0%-0,4%)

One-way ANOVA with the Dunnett post-test for multiple comparisons using untreated as the comparator

Conclusions and future perspectives

- ✓ *FLT3* expression is increased in *PAX5r* patients
- ✓ Gilteritinib is effective in *in-vitro*, *ex-vivo* and *in-vivo* in preclinical *PAX5r* models, and its anti-tumor activity is enhanced in the presence of standard chemotherapeutic drugs → **additive/synergistic effect**
- Study the expression of **FLT3 protein** and its active form, P-FLT3, in *PAX5r* patients
- Evaluate the efficacy and synergy of gilteritinib *in-vitro* and *in-vivo* in **PDX mouse models** of gilteritinib with other drugs, such as venetoclax

Acknowledgements

- Dr. Grazia Fazio
- Prof. Giovanni Cazzaniga
- Dr. Nicolò Peccatori
- Dr. Rebecca Rapisarda
- Dr. Domenico Gaspari
- Dr. Stefano Rebellato
- Dr. Chiara Palmi
- Dr. Michela Bardini
- Prof. Andrea Biondi



AIRC IG2023 to Grazia Fazio

